






## CLAIMS






We claim:

5 1. A compound of Formula (I), the racemic-diastereomeric mixtures, optical  
isomers, pharmaceutically-acceptable salts, prodrugs or biologically active  
metabolites thereof, selected from the group consisting of










*a /*  
*cont*




*a'*  
Cont




5





Q1  
Cont



*A'*  
cont





*A1  
cont*



where  $Z^{100}$  is or a group optionally substituted with R<sub>b</sub> selected from the group consisting of cycloalkyl, naphthyl, tetrahydronaphthyl,

benzothienyl, furanyl, thienyl, benzoxazolyl, benzothiazolyl,



 , thiazolyl, benzofuranyl, 2,3-dihydrobenzofuranyl, indolyl, isoxazolyl, tetrahydropyranyl, tetrahydrofuranyl, piperidinyl, pyrazolyl, pyrrolyl, oxazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, indolinyl, indazolyl, benzoisothiazolyl, pyrido-oxazolyl, pyrido-thiazolyl, pyrimido-oxazolyl, pyrimido-thiazolyl and benzimidazolyl;

10       $Z^{110}$  is a covalent bond, or an optionally substituted ( $C_1-C_6$ ) which is optionally substituted with one or more substituents selected from the group consisting of alkyl, CN, OH, halogen,  $NO_2$ , COOH, substituted or unsubstituted amino and substituted or unsubstituted phenyl;

15       $Z^{111}$  is a covalent bond, an optionally substituted ( $C_1-C_6$ ) or an optionally substituted

*A1*  
cont 5

- $(CH_2)_n$ -cycloalkyl- $(CH_2)_n$ -, where the optionally substituted groups are  
optionally substituted with one or more substituents selected from the group  
consisting of alkyl, CN, OH, halogen, NO<sub>2</sub>, COOH, substituted or unsubstituted  
amino and substituted or unsubstituted phenyl;

10 R<sub>a</sub> and R<sub>b</sub> each represent one or more substituents for each occurrence  
independently selected from the group consisting of hydrogen, halogen, -CN, -  
NO<sub>2</sub>, -C(O)OH, -C(O)H, -OH, -C(O)O-alkyl, substituted or unsubstituted  
carboxamido, tetrazolyl, trifluoromethylcarbonylamino,  
trifluoromethylsulfonamido, substituted or unsubstituted alkyl, substituted or  
unsubstituted alkoxy, substituted or unsubstituted aryl, substituted or  
unsubstituted alkenyl, substituted or unsubstituted aryloxy, substituted or  
unsubstituted heteroaryloxy, substituted or unsubstituted arylalkyl, substituted  
or unsubstituted alkynyl, substituted or unsubstituted amino, substituted or  
unsubstituted aminoalkyl, substituted or unsubstituted amido groups,  
substituted or unsubstituted heteroarylthio, substituted or unsubstituted arylthio,  
-Z<sup>105</sup>-C(O)N(R)<sub>2</sub>, -Z<sup>105</sup>-N(R)-C(O)-Z<sup>200</sup>, -Z<sup>105</sup>-N(R)-S(O)<sub>2</sub>-Z<sup>200</sup>, -Z<sup>105</sup>-N(R)-C(O)-  
N(R)-Z<sup>200</sup>, R<sub>c</sub> and CH<sub>2</sub>OR<sub>c</sub>;

15 where R<sub>c</sub> for each occurrence is independently hydrogen, substituted or  
unsubstituted alkyl, substituted or unsubstituted aryl, -CH<sub>2</sub>-NR<sub>d</sub>R<sub>e</sub>, -W-(CH<sub>2</sub>)<sub>t</sub>-  
NR<sub>d</sub>R<sub>e</sub>, -W-(CH<sub>2</sub>)<sub>t</sub>-Oalkyl, -W-(CH<sub>2</sub>)<sub>t</sub>-S-alkyl, or -W-(CH<sub>2</sub>)<sub>t</sub>-OH;  
20 Z<sup>105</sup> for each occurrence is independently a covalent bond or (C<sub>1</sub>-C<sub>6</sub>);  
Z<sup>200</sup> for each occurrence is independently a substituted or unsubstituted (C<sub>1</sub>-C<sub>6</sub>),  
substituted or unsubstituted phenyl or substituted or unsubstituted -(C<sub>1</sub>-C<sub>6</sub>)-  
phenyl;

25 R<sub>d</sub> and R<sub>e</sub> for each occurrence are independently H, alkyl, alkanoyl or  
SO<sub>2</sub>-alkyl; or R<sub>d</sub>, R<sub>e</sub> and the nitrogen atom to which they are attached  
together form a five- or six-membered heterocyclic ring; t for each  
occurrence is independently an integer from 2 to 6; W for each  
occurrence is independently a direct bond or O, S, S(O), S(O)<sub>2</sub>, or NR<sub>f</sub>,  
30 wherein R<sub>f</sub> for each occurrence is independently H or alkyl;  
or R<sub>i</sub> is a substituted or unsubstituted carbocyclic or heterocyclic ring fused

with

ring 2;

R<sub>3</sub> is hydrogen, hydroxy, substituted or unsubstituted alkyl or substituted or unsubstituted alkoxy;

5

A is -O-; -S-; -S(O)<sub>p</sub>-; -N(R)-; -N(C(O)OR)-; -N(C(O)R)-; -N(SO<sub>2</sub>R)-; -CH<sub>2</sub>O-; -CH<sub>2</sub>S-; -CH<sub>2</sub>N(R)-; -CH(NR)-; -CH<sub>2</sub>N(C(O)R))-; -CH<sub>2</sub>N(C(O)OR)-; -CH<sub>2</sub>N(SO<sub>2</sub>R)-; -CH(NHR)-; -CH(NHC(O)R)-; -CH(NHSO<sub>2</sub>R)-; -CH(NHC(O)OR)-; -CH(OC(O)R)-; -CH(OC(O)NHR); -CH=CH-; -C(=NOR)-; -C(O)-; -CH(OR)-; -C(O)N(R)-; -N(R)C(O)-; -N(R)S(O)<sub>p</sub>-; -QC(O)N(R)-; -N(R)-C(O)-(CH<sub>2</sub>)<sub>n</sub>-N(R)-; -N(R)C(O)O-; -N(R)-(CH<sub>2</sub>)<sub>n+1</sub>-C(O)-; -S(O)<sub>p</sub>N(R)-; -O-(CR<sub>2</sub>)<sub>n+1</sub>-C(O)-; -O-(CR<sub>2</sub>)<sub>n+1</sub>-O-; -N(C(O)R)S(O)<sub>p</sub>-; -N(R)S(O)<sub>p</sub>N(R)-; -N(R)-C(O)-(CH<sub>2</sub>)<sub>n</sub>-O-; -C(O)N(R)C(O)-; -S(O)<sub>p</sub>N(R)C(O)-; -OS(O)<sub>p</sub>N(R)-; -N(R)S(O)<sub>p</sub>O-; -N(R)S(O)<sub>p</sub>C(O)-; -SO<sub>p</sub>N(C(O)R)-; -N(R)SO<sub>p</sub>N(R)-; -C(O)O-; -N(R)P(OR<sub>g</sub>)O-; -N(R)P(OR<sub>g</sub>)-; -N(R)P(O)(OR<sub>g</sub>)O-; -N(R)P(O)(OR<sub>g</sub>)-; -N(C(O)R)P(OR<sub>g</sub>)O-; -N(C(O)R)P(OR<sub>g</sub>)-; -N(C(O)R)P(O)(OR<sub>g</sub>)O-, or -N(C(O)R)P(OR<sub>g</sub>)-;

10

where R for each occurrence is independently H, substituted or unsubstituted alkyl, substituted or unsubstituted arylalkyl or substituted or unsubstituted aryl;

15

R<sub>g</sub> for each occurrence is independently H, substituted or unsubstituted alkyl, substituted or unsubstituted arylalkyl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted aryl;

20

p is 1 or 2;

or in a phosphorus containing group, the nitrogen atom, the phosphorus atom, R and R<sub>g</sub> together form a five- or six-membered heterocyclic ring; or

A is NRSO<sub>2</sub> and R, R<sub>a</sub> and the nitrogen atom together form a substituted or unsubstituted five or-six-membered heterocyclic ring fused to ring 1;

25

R<sub>2</sub> is -Z<sup>101</sup>-Z<sup>102</sup>;

*Q*  
Cont

a  
cont  
5

$Z^{101}$  is a covalent bond,  $-(C_1-C_6)-$ ,  $-(C_1-C_6)-O-$ ,  $-(C_1-C_6)-C(O)-$ ,  $-(C_1-C_6)-C(O)O-$ ,  
 $-(C_1-C_6)-C(O)-NH-$ ,  $-(C_1-C_6)-C(O)-N((C_1-C_6))-$  or a substituted or  
unsubstituted phenyl group;

$Z^{102}$  is hydrogen, a substituted or unsubstituted alkyl group, a substituted or  
unsubstituted cycloalkyl group, a substituted or unsubstituted, saturated  
or unsaturated heterocyclic group, or a substituted or unsubstituted,  
saturated or unsaturated heterobicyclic group;

10 said substituted heterocyclic or substituted heterobicyclic group having one or  
more substituents each independently selected from the group consisting of  
hydroxyl, cyano, substituted or unsubstituted alkoxy, substituted or  
unsubstituted sulfonamido, substituted or unsubstituted ureido, substituted or  
unsubstituted carboxamido; substituted or unsubstituted amino, oxo, a saturated,  
unsaturated or aromatic, substituted or unsubstituted heterocyclic group  
comprising one or more nitrogen atoms, one or more oxygen atoms or a  
combination thereof;

15 wherein said nitrogen atoms are independently optionally substituted by  
a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or  
substituted or unsubstituted arylalkyl group; or

20  $R_2$  is of the formula B-E, wherein B is a substituted or unsubstituted cycloalkyl,  
substituted or unsubstituted azacycloalkyl, substituted or unsubstituted amino,  
substituted or unsubstituted aminoalkylsulfonyl, substituted or unsubstituted  
alkoxyalkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted  
aminoalkylcarbonyl, hydroxy, substituted or unsubstituted alkylene, substituted or  
unsubstituted aminoalkyl, substituted or unsubstituted alkylidenecarbonyl or substituted  
25 or unsubstituted aminoalkylcarbonyl group; and E is substituted or unsubstituted  
azacycloalkyl, substituted or unsubstituted azacycloalkylcarbonyl, substituted or  
unsubstituted azacycloalkylsulfonyl, substituted or unsubstituted azacycloalkylalkyl,  
substituted or unsubstituted heteroaryl, substituted or unsubstituted heteroarylcarbonyl,  
substituted or unsubstituted heteroarylsulfonyl, substituted or unsubstituted  
30 heteroarylalkyl, substituted or unsubstituted azacycloalkylcarbonylamino, substituted or  
unsubstituted heteroarylcarbonylamino or substituted or unsubstituted aryl;

*A*  
*cont*

a is 1 and D<sub>1</sub>, G<sub>1</sub>, J<sub>1</sub>, L<sub>1</sub> and M<sub>1</sub> are each independently selected from the group consisting of CR<sub>a</sub> and N, provided that at least two of D<sub>1</sub>, G<sub>1</sub>, J<sub>1</sub>, L<sub>1</sub> and M<sub>1</sub> are CR<sub>a</sub>; or

5        a is 0, and one of D<sub>1</sub>, G<sub>1</sub>, L<sub>1</sub> and M<sub>1</sub> is NR<sub>a</sub>, one of D<sub>1</sub>, G<sub>1</sub>, L<sub>1</sub> and M<sub>1</sub> is CR<sub>a</sub> and the remainder are independently selected from the group consisting of CR<sub>a</sub> and N, wherein R<sub>a</sub> is as defined above;

b is 1 and D<sub>2</sub>, G<sub>2</sub>, J<sub>2</sub>, L<sub>2</sub> and M<sub>2</sub> are each independently selected from the group consisting of CR<sub>a</sub> and N, provided that at least two of D<sub>2</sub>, G<sub>2</sub>, J<sub>2</sub>, L<sub>2</sub> and M<sub>2</sub> are CR<sub>a</sub>; or


10      b is 0, and one of D<sub>2</sub>, G<sub>2</sub>, L<sub>2</sub> and M<sub>2</sub> is NR<sub>a</sub>, one of D<sub>2</sub>, G<sub>2</sub>, L<sub>2</sub> and M<sub>2</sub> is CR<sub>a</sub> and the remainder are independently selected from the group consisting of CR<sub>a</sub> and N, wherein R<sub>a</sub> is as defined above; and

n for each occurrence is independently an integer from 0 to 6.

15      2. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>1</sub> for each occurrence is independently selected from the group consisting of F, Cl, Br, I, CH<sub>3</sub>, NO<sub>2</sub>, OCF<sub>3</sub>, OCH<sub>3</sub>, CN, CO<sub>2</sub>CH<sub>3</sub>, CF<sub>3</sub>, -CH<sub>2</sub>NR<sub>d</sub>R<sub>e</sub>, t-butyl, pyridyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted benzyl, substituted or unsubstituted benzenesulfonyl, substituted or unsubstituted phenoxy, substituted or unsubstituted phenyl, substituted or unsubstituted amino, carboxyl, substituted or unsubstituted tetrazolyl, and substituted or unsubstituted styryl.

20      3. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>a</sub> for each occurrence is independently selected from the group consisting of F, Cl, Br, I, CH<sub>3</sub>, NO<sub>2</sub>, OCF<sub>3</sub>, OCH<sub>3</sub>, CN, CO<sub>2</sub>CH<sub>3</sub>, CF<sub>3</sub>, t-butyl, pyridyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted benzyl, substituted or unsubstituted benzenesulfonyl, substituted or unsubstituted phenoxy, substituted or unsubstituted phenyl, substituted or unsubstituted amino, carboxyl, substituted or unsubstituted tetrazolyl, and substituted or unsubstituted styryl.

4. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula




5

wherein n is 1, 2 or 3.

5. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

10



15

wherein m is 0, 1, 2 or 3 and


R<sub>g</sub> is H or -(CH<sub>2</sub>)<sub>p</sub>N(R<sub>4</sub>)R<sub>5</sub>, wherein p is an integer from 2 to 6 and R<sub>4</sub> and R<sub>5</sub> are each, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted moiety selected from the group consisting of alkyl, alkoxy, amino, aryl, heteroaryl and heterocycloalkyl group or R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom to which they are attached together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group.

20

25

6. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

30



wherein m is 0, 1, 2 or 3

a and b are each, independently, an integer from 0 to 6;

Q is -OR<sub>6</sub> or -NR<sub>4</sub>R<sub>5</sub>;

each R<sub>4</sub> and R<sub>5</sub> is, independently, H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>-, wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, amino, aryl, heteroaryl or heterocycloalkyl group or


10 R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom to which they are attached together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or heterobicyclic group; and

R<sub>6</sub> is hydrogen or a substituted or unsubstituted alkyl group.

15

7. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

5



wherein n is 1, 2 or 3; and

R<sub>4</sub> is H, azabicycloalkyl or Y-Z, wherein Y is selected

10


from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>-; wherein q is an integer 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted amino, aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted

15

heterocycloalkyl group.

8. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

20



wherein

25

m is 0, 1, 2 or 3;

R<sub>5</sub> is H, azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of a covalent bond, -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, -(CH<sub>2</sub>)<sub>q</sub>C(O)-, -C(O)(CH<sub>2</sub>)<sub>q</sub>- and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>, where the alkyl portion of -(CH<sub>2</sub>)<sub>q</sub>-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, -(CH<sub>2</sub>)<sub>q</sub>C(O)-, -C(O)(CH<sub>2</sub>)<sub>q</sub>- and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>, is optionally substituted by a halogen, hydroxy or an alkyl group;

30

wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted alkoxy, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or Y and Z together are a natural or unnatural amino acid, which may be mono- or di-alkylated at the amine nitrogen; and

5

R<sub>6</sub> represents one or more substituents each independently selected from the group consisting of hydrogen, hydroxy, oxo, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl, substituted or unsubstituted arylcarbonyl, substituted or unsubstituted heterocyclcarbonyl, substituted or unsubstituted aminoalkyl and substituted or unsubstituted arylalkyl; provided that the carbon atoms adjacent to the nitrogen atom are not substituted by a hydroxy group.


10

15

9. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

20

25




30

wherein R<sub>4</sub> is H, substituted or unsubstituted alkyl, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein q is an integer from 0 to 6, and r is 0, 1 or 2; and Z is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl

or substituted or unsubstituted heterocycloalkyl.

10. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

5



10

wherein

m is an integer from 1 to 6; and

15


R<sub>4</sub> and R<sub>5</sub> are each, independently, H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

20

R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom to which they are attached together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or substituted or unsubstituted heterobicyclic group.

11. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

25



30

wherein

n is an integer from 0 to 4;

r is 0 and m is an integer from 1 to 6; or

5 r is 1 and m is an integer from 0 to 6;

Q is -OR<sub>6</sub> or -NR<sub>4</sub>R<sub>5</sub>;


each R<sub>4</sub> and R<sub>5</sub> is, independently, H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>-

10 ; q is an integer from 0 to 6; and r is 0, 1 or 2; and Z is a substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

15 R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom to which they are attached together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group; and R<sub>6</sub> is hydrogen or a substituted or unsubstituted alkyl group.

12. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

20



25

n is an integer from 0 to 4;

m is an integer from 0 to 6;

30

R<sub>4</sub> is H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is

selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -


SO<sub>2</sub>NH-, -CONH-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>-; wherein q is an

integer from 0 to 6; and r is 0, 1 or 2; and Z is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; and R<sub>6</sub> is hydrogen or a substituted or unsubstituted alkyl group.

5

13. The compound of Claim 10 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocyclic group of the formula

10



wherein

15

R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub> and R<sub>14</sub> are each, independently, lower alkyl or hydrogen; or at least one pair of substituents R<sub>7</sub> and R<sub>8</sub>; R<sub>9</sub> and R<sub>10</sub>; R<sub>11</sub> and R<sub>12</sub>; or R<sub>13</sub> and R<sub>14</sub> together are an oxygen atom; or at least one of R<sub>7</sub> and R<sub>8</sub> is cyano, CONHR<sub>15</sub>, COOR<sub>15</sub>, CH<sub>2</sub>OR<sub>15</sub> or CH<sub>2</sub>NR<sub>15</sub>(R<sub>16</sub>), wherein R<sub>15</sub> and R<sub>16</sub> are each, independently, H, azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-,-S(O)<sub>2</sub>-,-C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or R<sub>15</sub>, R<sub>16</sub> and the nitrogen atom together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or a substituted or unsubstituted heterobicyclic group;

20

X is O, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, CHOR<sub>17</sub> or NR<sub>17</sub>, wherein R<sub>17</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, -C(NH)NH<sub>2</sub>, -C(O)R<sub>17</sub>, or -C(O)OR<sub>18</sub>, wherein R<sub>18</sub> is

25

hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; and


30

t is 0 or 1.

14. The compound of Claim 10 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocycle of the formula

5

10



wherein

R<sub>19</sub> and R<sub>20</sub> are each, independently, hydrogen or lower alkyl; or R<sub>19</sub> and R<sub>20</sub> together are an oxygen atom;

15

R<sub>21</sub> and R<sub>22</sub> are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

20

R<sub>21</sub>, R<sub>22</sub> and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group;

25


m is an integer from 1 to 6; and

n is an integer from 0 to 6.

30

15. The compound of Claim 10 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocyclic group of the formula

5



wherein

m is an integer from 1 to 6; and

10 R<sub>23</sub> is CH<sub>2</sub>OH, NRR', C(O)NRR' or COOR, wherein R and R' are each, independently, hydrogen or substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl.


15. The compound of Claim 10 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocyclic group of the formula

20

wherein R<sub>24</sub> is substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl, carboxyl, cyano, C(O)OR<sub>25</sub>, CH<sub>2</sub>OR<sub>25</sub>, CH<sub>2</sub>NR<sub>26</sub>R<sub>27</sub> or C(O)NHR<sub>26</sub>, wherein R<sub>25</sub> is substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, substituted or unsubstituted heterocyclic or substituted or unsubstituted heterocycloaryl; and R<sub>26</sub> and R<sub>27</sub> are each, independently, H,


25 substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-,-S(O)<sub>2</sub>-,-C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or

30



unsubstituted heterocycloalkyl; or R<sub>26</sub>, R<sub>27</sub> and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group.


17. The compound of Claim 10 wherein at least one of R<sub>4</sub> and R<sub>5</sub> is of the formula  
 5 Y-Z, wherein Z is of the formula



10 wherein  
 T is C(O), S, SO, SO<sub>2</sub>, CHOR or NR, wherein R is hydrogen or a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl group; and  
 n is 0, 1 or 2.

15 18. The compound of Claim 10 wherein at least one of R<sub>4</sub> and R<sub>5</sub> is of the formula  
 Y-Z, wherein Z is of the formula -N(R<sub>28</sub>)R<sub>29</sub>, wherein R<sub>28</sub> and R<sub>29</sub> are each, independently, substituted or unsubstituted carboxyalkyl, substituted or unsubstituted alkoxy carbonylalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted cyanoalkyl; or  
 20 R<sub>28</sub> and R<sub>29</sub>, together with the nitrogen atom, form a five- or six-membered substituted or unsubstituted heterocyclic group.

25 19. The compound of Claim 11 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocycle of the formula



wherein

$R_7, R_8, R_9, R_{10}, R_{11}, R_{12}, R_{13}$  and  $R_{14}$  are each, independently, lower alkyl or hydrogen; or at least one pair of substituents  $R_7$  and  $R_8$ ;  $R_9$  and  $R_{10}$ ;  $R_{11}$  and  $R_{12}$ ;

5           or  $R_{13}$  and  $R_{14}$  together are an oxygen atom; or at least one of  $R_7$  and  $R_9$  is cyano,  $CONHR_{15}$ ,  $COOR_{15}$ ,  $CH_2OR_{15}$  or  $CH_2NR_{15}(R_{16})$ , wherein  $R_{15}$  and  $R_{16}$  are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of  $-C(O)-$ ,  $-(CH_2)_p-$ ,  $-S(O)_2-$ ,  $-C(O)O-$ ,  $-SO_2NH-$ ,  $-CONH-$ ,  $(CH_2)_qO-$ ,  $-(CH_2)_qNH-$ , and  $-(CH_2)_qS(O)_r-$ ; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is

10          substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or  $R_{15}, R_{16}$  and the nitrogen atom together form a 3, 4, 5, 6 or 7-membered, substituted or unsubstituted heterocyclic or

15          heterobicyclic group;


X is O, S, SO,  $SO_2$ ,  $CH_2$ ,  $CHOR_{17}$  or  $NR_{17}$ , wherein  $R_{17}$  is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or

unsubstituted arylalkyl,  $-C(NH)NH_2$ ,  $-C(O)R_{18}$ , or  $-C(O)OR_{18}$ , wherein  $R_{18}$  is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or

20          substituted or unsubstituted arylalkyl; and

t is 0 or 1.

25          20. The compound of Claim 11 wherein  $R_4, R_5$  and the nitrogen atom together form a heterocycle of the formula



30

wherein


R<sub>19</sub> and R<sub>20</sub> are each, independently, hydrogen or lower alkyl; or R<sub>19</sub> and R<sub>20</sub> together are an oxygen atom;

5           R<sub>21</sub> and R<sub>22</sub> are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>-; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

10          R<sub>21</sub>, R<sub>22</sub> and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group;

15          m is an integer from 1 to 6; and  
n is an integer from 0 to 6.

21. The compound of Claim 11 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocyclic group of the formula




25          wherein

m is an integer from 1 to 6; and

R<sub>23</sub> is CH<sub>2</sub>OH, NRR', C(O)NRR' or COOR, wherein R is hydrogen or a substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl group.

22. The compound of Claim 11 wherein R<sub>4</sub>, R<sub>5</sub> and the nitrogen atom together form a heterocyclic group of the formula

5



10


15

20

wherein R<sub>24</sub> is substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl, carboxyl, cyano, C(O)OR<sub>25</sub>, CH<sub>2</sub>OR<sub>25</sub>, CH<sub>2</sub>NR<sub>26</sub>R<sub>27</sub> or C(O)NHR<sub>26</sub>, wherein R<sub>25</sub> is substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, substituted or unsubstituted heterocyclic or substituted or unsubstituted heterocycloaryl group; and R<sub>26</sub> and R<sub>27</sub> are each, independently, H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-,-S(O)<sub>2</sub>-,-C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or R<sub>26</sub>, R<sub>27</sub> and the nitrogen atom together form a 3, 4, 5 or 6-membered, substituted or unsubstituted heterocyclic group.

23. The compound of Claim 11 wherein at least one of R<sub>4</sub> and R<sub>5</sub> is of the formula Y-Z, wherein Z is of the formula

25



30

wherein

g is 0 or 1;

T is C(O), O, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, CHOR<sub>17</sub> or NR<sub>17</sub>, wherein R<sub>17</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, -C(NH)NH<sub>2</sub>, -C(O)R<sub>18</sub>, or -C(O)OR<sub>18</sub>, wherein R<sub>18</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; and

R<sub>32</sub> is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl.

5

10

15


20

25

24. The compound of Claim 11 wherein at least one of R<sub>4</sub> and R<sub>5</sub> is of the formula Y-Z, wherein Z is of the formula -N(R<sub>28</sub>)R<sub>29</sub>, wherein R<sub>28</sub> and R<sub>29</sub> are each, independently, substituted or unsubstituted carboxyalkyl, substituted or unsubstituted alkoxycarbonylalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted alkylsulfonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted cyanoalkyl; or R<sub>28</sub> and R<sub>29</sub>, together with the nitrogen atom, form a five- or six-membered substituted or unsubstituted heterocyclic group.
25. The compound of Claim 8 wherein R<sub>5</sub> is Y-Z, wherein Z is of the formula N(R<sub>30</sub>)R<sub>31</sub>, wherein R<sub>30</sub> and R<sub>31</sub> are each, independently, hydrogen, alkyl, alkoxycarbonyl, alkoxyalkyl, hydroxyalkyl, aminocarbonyl, cyano, alkylcarbonyl or arylalkyl.

26. The compound of Claim 8 wherein R<sub>5</sub> is Y-Z, wherein Z is of the formula

5



wherein


each X is, independently, CH or N; and

10 R<sub>32</sub> is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.

15

27. The compound of Claim 8 wherein R<sub>5</sub> is Y-Z, wherein Z is of the formula

20



wherein

g is 0 or 1;

25


T is O, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, CHOR<sub>17</sub> or NR<sub>17</sub>, wherein R<sub>17</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted arylalkyl, C(O)NH<sub>2</sub>, -C(NH)NH<sub>2</sub>, -C(O)R<sub>17</sub>, or -C(O)OR<sub>18</sub>, wherein R<sub>18</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; and

30

R<sub>32</sub> is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl,

substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.

5 28. The compound of Claim 8 wherein R<sub>5</sub> is Y-Z, wherein Z is of the formula




10 wherein

g is 0, 1 or 2; and

15 R<sub>32</sub> is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl, substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.

29. The compound of Claim 8 wherein R<sub>5</sub> is Y-Z, wherein Z is of the formula

20




wherein

25 T is C(O), O, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, CHOR<sub>17</sub> or NR<sub>17</sub>, wherein R<sub>17</sub> is hydrogen, substituted or unsubstituted alkyl, aryl, arylalkyl, -C(NH)NH<sub>2</sub>, -C(O)R<sub>18</sub>, or -C(O)OR<sub>18</sub>, wherein R<sub>18</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl or substituted or unsubstituted arylalkyl; g is 0 or 1; and

30 R<sub>32</sub> is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl,

substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl group.

5 30. The compound of Claim 8 wherein R<sub>5</sub> is Y-Z, wherein Z is of the formula



10

wherein

R<sub>32</sub> is hydrogen, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl,

15

substituted or unsubstituted hydroxyalkyl, substituted or unsubstituted aminocarbonyl, alkylcarbonyl, substituted or unsubstituted thioalkoxy or substituted or unsubstituted arylalkyl; and


20

R<sub>33</sub> is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxycarbonyl, substituted or unsubstituted alkoxyalkyl,

substituted or unsubstituted aminocarbonyl, perhaloalkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkylcarbonyl or substituted or unsubstituted arylalkyl.

31. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula

25



30

wherein


m is 0 or 1;

R<sub>34</sub>, R<sub>35</sub>, R<sub>36</sub>, R<sub>37</sub>, R<sub>38</sub>, R<sub>39</sub>, R<sub>40</sub> and R<sub>41</sub> are each, independently, methyl or hydrogen; or at least one pair of substituents R<sub>34</sub> and R<sub>35</sub>; R<sub>36</sub> and R<sub>37</sub>; R<sub>38</sub> and R<sub>39</sub>; or R<sub>40</sub> and R<sub>41</sub> together are an oxygen atom; and

5 R<sub>42</sub> is H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-,-S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and Z is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl group; or

10 R<sub>42</sub> is of the formula

15



wherein

20

u is 0 or 1;

R<sub>43</sub>, R<sub>44</sub>, R<sub>45</sub>, R<sub>46</sub>, R<sub>47</sub>, R<sub>48</sub>, R<sub>49</sub> and R<sub>50</sub> are each, independently, methyl or hydrogen;

or at least one pair of substituents R<sub>43</sub> and R<sub>44</sub>; R<sub>45</sub> and R<sub>46</sub>; R<sub>47</sub> and R<sub>48</sub>;

or R<sub>49</sub> and R<sub>50</sub> together are an oxygen atom; and


25

R<sub>51</sub> is H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-,-S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or

30

unsubstituted heterocycloalkyl.

32. The compound of Claim 1 wherein R<sub>3</sub> is H; R<sub>2</sub> is of the formula




wherein

h, i, j, k and l are independently 0 or 1;

R<sub>52</sub>, R<sub>53</sub>, R<sub>54</sub>, R<sub>55</sub>, R<sub>56</sub>, R<sub>57</sub>, R<sub>58</sub>, R<sub>59</sub>, R<sub>g</sub> and R<sub>h</sub> are each, independently, methyl or hydrogen; or at least one pair of substituents R<sub>52</sub> and R<sub>53</sub>; R<sub>54</sub> and R<sub>55</sub>; R<sub>56</sub> and R<sub>57</sub>; or R<sub>58</sub> and R<sub>59</sub> together are an oxygen atom; and

15 R<sub>60</sub> is H, substituted or unsubstituted azabicycloalkyl or Y-Z, wherein Y is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-,-S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and-(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and Z is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl; or

20 R<sub>60</sub> is of the formula



wherein

30 v is 0 or 1;

$R_{61}$ ,  $R_{62}$ ,  $R_{63}$ ,  $R_{64}$ ,  $R_{65}$ ,  $R_{66}$ ,  $R_{67}$  and  $R_{68}$  are each, independently, lower alkyl or hydrogen; or at least one pair of substituents  $R_{61}$  and  $R_{62}$ ;  $R_{63}$  and  $R_{64}$ ;  $R_{65}$  and  $R_{66}$ ; and  $R_{67}$  and  $R_{68}$  together are an oxygen atom; and  
5            $R_{69}$  is H, substituted or unsubstituted azabicycloalkyl or V-L, wherein V is selected from the group consisting of -C(O)-, -(CH<sub>2</sub>)<sub>p</sub>-, -S(O)<sub>2</sub>-, -C(O)O-, -SO<sub>2</sub>NH-, -CONH-, (CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>NH-, and -(CH<sub>2</sub>)<sub>q</sub>S(O)<sub>r</sub>-; wherein p is an integer from 0 to 6, q is an integer from 0 to 6, and r is 0, 1 or 2; and L is substituted or unsubstituted alkyl, substituted or unsubstituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl or substituted or unsubstituted heterocycloalkyl.  
10

*Sub C2*  
33. A method of inhibiting one or more protein kinase activity in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.

15           34. The method of Claim 33 wherein said protein kinase is selected from the group consisting of KDR, FGFR-1, PDGFR $\beta$ , PDGFR $\alpha$ , IGF-1R, c-Met, Flt-1, Flt-4, TIE-2, TIE-1, Lck, Src, fyn, Lyn, Blk, hck, fgr and yes.

*Sub C3*  
20           35. A method of affecting hyperproliferative disorders in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.

25           36. A method of affecting angiogenesis in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.

30           37. The method of Claim 33 wherein the protein kinase is a protein serine/threonine kinase or a protein tyrosine kinase.

Sub  
C7

38. A method of treating one or more ulcers in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient.

5

39. The method of Claim 38 wherein the ulcer or ulcers are caused by a bacterial or fungal infection; or the ulcer or ulcers are Mooren ulcers; or the ulcer or ulcers are a symptom of ulcerative colitis.

10 40. A method of treating a condition in a patient comprising administering a therapeutically effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolites thereof to said patient, wherein said condition is an ocular condition, a cardiovascular condition, a cancer, Crow-Fukase (POEMS) syndrome, a diabetic condition, sickle cell anaemia, chronic inflammation, systemic lupus, glomerulonephritis, synovitis, inflammatory bowel disease, Crohn's disease, glomerulonephritis, rheumatoid arthritis, osteoarthritis, multiple sclerosis, graft rejection, Lyme disease, sepsis, von Hippel Lindau disease, pemphigoid, psoriasis, Paget's disease, polycystic kidney disease, fibrosis, sarcoidosis, cirrhosis, thyroiditis, hyperviscosity syndrome, Osler-Weber-Rendu disease, chronic occlusive pulmonary disease, asthma or edema following burns, trauma, radiation, stroke, hypoxia, ischemia, ovarian hyperstimulation syndrome, preeclampsia, menometrorrhagia, endometriosis, or infection by Herpes simplex, Herpes Zoster, human immunodeficiency virus, parapoxvirus, protozoa or toxoplasmosis.

15

20 41. The method of Claim 40 wherein the ocular condition is ocular or macular edema, ocular neovascular disease, scleritis, radial keratotomy, uveitis, vitritis, myopia, optic pits, chronic retinal detachment, post-laser treatment complications, conjunctivitis, Stargardt's disease, Eales disease, retinopathy or macular degeneration.

25

Sub  
CS

42. The method of Claim 40 wherein the cardiovascular condition is atherosclerosis, restenosis, ischemia/reperfusion injury, vascular occlusion or carotid obstructive disease.

5

43. The method of Claim 40 wherein the cancer is a solid tumor, a sarcoma, fibrosarcoma, osteoma, melanoma, retinoblastoma, a rhabdomyosarcoma, glioblastoma, neuroblastoma, teratocarcinoma, an hematopoietic malignancy, Kaposi's sarcoma, Hodgkin's disease, lymphoma, myeloma, leukemia or malignant ascites.

10

44. The method of Claim 40 wherein the diabetic condition is insulin-dependent diabetes mellitus glaucoma, diabetic retinopathy or microangiopathy.

15

45. A method of decreasing fertility in a patient, said method comprising the step of administering to the patient an effective amount of a compound of Claim 1 or a physiologically acceptable salt, prodrug or biologically active metabolite thereof.

Sub  
C 6

20

46. The method of Claim 36 wherein the compound or a physiologically acceptable salt, prodrug or biologically active metabolite thereof is administered in an amount effective to promote angiogenesis or vasculogenesis.

25

47. The method of Claim 34 wherein the protein kinase is Tie-2.

Sub  
C 7

48. The method of Claim 46 wherein the compound of Formula I, or physiologically acceptable salt, prodrug or biologically active metabolite thereof, is administered in combination with a pro-angiogenic growth factor.


49. The method of Claim 48 wherein the pro-angiogenic growth factor is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, VEGF-E, HGF, FGF-1, FGF-2, derivatives thereof and antiiodotypic antibodies.


5 50. The method of Claim 46 wherein the patient is suffering from anemia, ischemia, infarct, transplant rejection, a wound, gangrene or necrosis.


10 51. The method of Claim 33 wherein the protein kinase activity is involved in T cell activation, B cell activation, mast cell degranulation, monocyte activation, the potentiation of an inflammatory response or a combination thereof.

15 52. A compound according to Claim 1, wherein  $R_3$  is H;  $R_2$  is  $-Z^{101}-Z^{102}$  where  $Z^{101}$  is a covalent bond,  $-(C_1-C_6)-$ ,  $-(C_1-C_6)-O-$ ,  $-(C_1-C_6)-C(O)-$ ,  $-(C_1-C_6)-C(O)O-$ ,  $-(C_1-C_6)-C(O)-NH-$ ,  $-(C_1-C_6)-C(O)-N((C_1-C_6))-$  or a substituted phenyl group; and  $Z^{102}$  is hydrogen, a substituted or unsubstituted alkyl group or a substituted or unsubstituted, saturated or unsaturated heterocyclic group.

20 53. A compound according to Claim 52, wherein  $Z^{101}$  is selected from the group consisting of  $-CH_2-C(O)O-$ ,  $-CH_2-C(O)-$ ,  $-CH_2-C(O)-NH-$ ,  $-CH_2-C(O)-N(Me)-$ ,  $-CH(Me)-C(O)O-$ ,  $-(CH_2)_3-C(O)O-$ ,  $-CH(Me)-C(O)-NH-$ , and  $-(CH_2)_3-C(O)-NH-$ ;  $Z^{102}$  is selected from the group consisting of hydrogen, methyl, ethyl, N,N-dimethylaminoethyl, N,N-diethylaminoethyl, 2-phenyl-2-hydroxyethyl, morpholino, piperazinyl, N-methylpiperazinyl and 2-hydroxymethylpyrrolidinyl.

25 54. A compound according to Claim 53, wherein  $R_1$  is  ,





or 

where


$Z^{100}$  is a substituted or unsubstituted benzoxazolyl or a substituted or unsubstituted benzthiazolyl.

55. A compound according to Claim 8, 9, 10 or 53, wherein  $R_1$  is



5


or




where there is only one  $R_a$  and it is H or F.

56. A compound according to Claim 52, wherein  $Z^{101}$  is a covalent bond; and  $Z^{102}$  is  
10 an optionally substituted pyridyl.

15 57. A compound according to Claim 56, wherein  $R_1$  is



58. A compound according to Claim 1, wherein  $R_3$  is H;  
 $R_2$  is cyclopentyl; and



59. A compound according to Claim 58, wherein  
5       $Z^{110}$  is hydrogen;  
A is O; and  $Z^{100}$  is optionally substituted phenyl, furanyl or thienyl, where  $Z^{100}$  is  
optionally substituted with one or more substituents each independently selected  
from the group consisting of F, COOH, NO<sub>2</sub>, OMe, -COOMe, OCF<sub>3</sub> and CF<sub>3</sub>.

60. A compound according to Claim 58, wherein  
10      $Z^{110}$  is hydrogen;  
A is -O-, -O-(CR<sub>2</sub>)<sub>n</sub>-C(O)- or -O-(CR<sub>2</sub>)<sub>n</sub>-O-;  
n for each occurrence is 0 to 3;  
 $Z^{100}$  is an optionally substituted group selected from the group consisting of  
cyclohexyl, phenyl, tetrahydropyranyl, tetrahydrofuryl, isoxazolyl and  
15     piperidinyl; where  $Z^{100}$  is optionally substituted with one or more substituents  
selected from the group consisting of alkyl, alkoxy, halo, hydroxy and  
alkoxycarbonyl.

61. A compound according to Claim 58, wherein R<sup>2</sup> is an optionally substituted  
20     group selected from the group consisting of cyclobutyl and cyclohexyl.


62. A compound according to Claim 61, wherein R<sup>2</sup> is optionally substituted with  
one or more substituents selected from the group consisting of hydroxy, alkyl,  
hydroxyalkyl, carboxyalkyl and phenylalkoxyalkyl.  
25

63. A compound according to Claim 62, wherein R<sub>1</sub> is 4-phenoxyphenyl.

64. A compound according to Claim 6 wherein


m is 2; a is 0; R<sub>6</sub> is H; b is 1 or 2; and R<sub>4</sub> and R<sub>5</sub> are each hydrogen.

65. A compound according to Claim 8, wherein  
m is 0, 1 or 2; R<sub>6</sub> is hydrogen; R<sub>5</sub> is H or Y-Z;  
5 where Y is a covalent bond, -C(O)-, -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>q</sub>-, -(CH<sub>2</sub>)<sub>q</sub>C(O)- or -C(O)(CH<sub>2</sub>)<sub>q</sub>-, where the alkyl portion of -(CH<sub>2</sub>)<sub>q</sub>O-, -(CH<sub>2</sub>)<sub>p</sub>-, -(CH<sub>2</sub>)<sub>q</sub>C(O)- and -C(O)(CH<sub>2</sub>)<sub>q</sub>- is optionally substituted by a halogen, hydroxy or an alkyl group; and  
Z is hydrogen, alkyl, optionally substituted alkyl, alkoxyalkyl, optionally substituted heterocycloalkyl, optionally substituted heteroaryl, or optionally substituted amino.  
10

66. A compound according to Claim 65, wherein  
Z is hydrogen, methyl, ethyl, hydroxymethyl, methoxyethyl, N-methyl-piperidinyl, (t-butoxycarbonyl)(hydroxy)-piperidinyl, hydroxypiperidinyl,  
15 (hydroxymethyl)piperdinyl, (hydroxy)(methyl)-piperidinyl, morpholino, (methoxyethyl)piperizinyl, methylpiperizinyl, 4-piperidinylpiperidinyl, imidazolyl, methylimidazolyl, N-methylamino, N,N-dimethylamino, N-isopropylamino, N,N-diethylamino, 2,3-dihydroxypropylamino, 2-  
20 hydroxyethylamino, 3-hydroxypropylamino, methoxyethylamino,  
25 ethoxycarbonylmethylamino, phenylmethylamino, N-methyl-N-methoxyamino,  
hydroxyethylamino, 3-hydroxypropylamino, methoxyethylamino,  
ethoxycarbonylmethylamino, phenylmethylamino, N-methyl-N-methoxyamino,  
 , furanymethylamino, piperidinylethylamino, N-(2-N,N-dimethylaminoethyl)-N-methylamino, 2-N,N-dimethylaminoethylamino, N-methyl-N-(N-methylpiperidin-4-yl)amino, 2-morpholino-ethylamino, 3-morpholino-propylamino, 3-imidazolylpropylamino, or 3-(2-

oxopyrrolidinyl)propylamino.


67. A compound according to Claim 8, wherein m is 2; R<sub>5</sub> is Y-Z; Y is -C(O)-; and



5

68. A compound according to Claim 9, wherein

R<sub>4</sub> is hydrogen or methyl;



A is selected from the group consisting of O, -N(R)- and -N(R)C(O)-;

10 Z<sup>111</sup> is -(CH<sub>2</sub>)<sub>n</sub>-cycloalkyl-(CH<sub>2</sub>)<sub>n</sub>-;

R is hydrogen or alkyl;

n is 0 to 5;

R<sub>a</sub> is one or more substituents each independently selected from the group consisting of H, OH, F, Cl, methyl and methoxy; and

15 R<sub>b</sub> is one or more substituents each independently selected from the group consisting of H, CN, F, CF<sub>3</sub>, OCF<sub>3</sub>, methyl, methoxy and an optionally substituted amino group;


where said amino group is optionally substituted with one or two groups each independently selected from the group consisting of alkyl,

20 alkoxyalkyl, phenyl, substituted phenyl, and optionally substituted heteroaryl.

69. A compound according to Claim 68, wherein R<sub>b</sub> is 4-methylphenylthio or 2-pyridinylthio.

25

70. A compound according to Claim 9, wherein




where  $\text{Z}^{100}$  is selected from the group consisting of benzo[b]thiophene, furanyl and thiophene.

5


71. A compound according to Claim 9C, wherein  $\text{R}_a$  is alkoxy; A is  $-\text{NH}-\text{C}(\text{O})-$ ; and there is a covalent bond between A and  $\text{Z}^{100}$ .

72. A compound according to Claims 1, 8 or 9, wherein



10

A is selected from the group consisting of  $-\text{N}(\text{R})-\text{C}(\text{O})-\text{N}(\text{R})-$ ,  $-(\text{CH}_2)_n-$   $\text{N}(\text{R})\text{C}(\text{O})\text{N}(\text{R})-$ ,  $-\text{N}(\text{R})-$  and  $-\text{N}(\text{R})-\text{SO}_2-$ ; R is hydrogen or alkyl;



15

X is S, O or NR where R for each occurrence is independently H or Me;  
 $\text{R}_a$  is one or more substituents each independently selected from the group consisting of H and F; and  
 $\text{R}_b$  is one or more substituents each independently selected from the group consisting of H, F, Cl, Br,  $\text{NO}_2$ ,  $\text{CF}_3$ , alkyl, alkoxy and alkoxycarbonyl.


20

73. A compound according to Claim 72, wherein  
 $\text{R}_4$  is methyl; m is 1, 2 or 3;  $\text{R}_5$  is Y-Z, where Y is  $-\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{O})-$  or  $-\text{C}(\text{O})-(\text{CH}_2)_p-$ ; and Z is aminoalkyl, N-alkylamino, N,N-dialkylamino or

hydroxyalkylaminoalkyl.

74. A compound according to Claim 9, wherein


$R_4$  is methyl;  $R_1$  is



5

where  $n$  is 0 to 3;  $Z^{100}$  is an optionally substituted group selected from the group consisting of indolyl, indenyl, methylindenyl, methylindolyl, dimethylaminophenyl, phenyl, cyclohexyl and benzofuranyl.

10 75. A compound according to claim 9, wherein



$R_1$  is

$Z^{100}$  is an optionally substituted group selected from the group consisting of phenyl, imidazolyl, indolyl, furanyl, benzofuranyl and 2,3-dihydrobenzofuranyl;

15

where  $Z^{100}$  is optionally substituted with one or more substituents each independently selected from the group consisting of F, Cl, CN, optionally substituted alkyl, -O-(optionally substituted alkyl), -COOH, - $Z^{105}$ -C(O)N(R)<sub>2</sub>, - $Z^{105}$ -N(R)-C(O)- $Z^{200}$ , - $Z^{105}$ -N(R)-S(O)<sub>2</sub>- $Z^{200}$ , and - $Z^{105}$ -N(R)-C(O)-N(R)- $Z^{200}$ ;

$Z^{105}$  is a covalent bond or (C<sub>1</sub>-C<sub>6</sub>);

20

$Z^{200}$  is an optionally substituted group selected from group consisting of (C<sub>1</sub>-C<sub>6</sub>), phenyl and -(C<sub>1</sub>-C<sub>6</sub>)-phenyl;


$Z^{110}$  and  $Z^{111}$  are each independently a covalent bond or (C<sub>1</sub>-C<sub>3</sub>) group optionally substituted with alkyl, hydroxy, COOH, CN or phenyl; and

A is O, -N(R)-C(O)-N(R)-, -N(R)-C(O)-O-, -N(R)- or -N(R)-C(O)-, where R is H or alkyl.

76. A compound according to Claim 75, wherein R<sub>4</sub> is methyl.

5

77. A compound according to Claim 8, 9 or 10, wherein




10


78. A compound according to Claim 77, wherein R<sub>4</sub> is methyl; A is -NH-; there is only one R<sub>a</sub> and it is H or F; and Z<sup>100</sup> is optionally substituted with one or more substituents each independently selected from the group consisting of alkyl, halo, CF<sub>3</sub>, and alkoxy.

15

79. A compound according to Claim 9, wherein



20



where Z<sup>100</sup> is optionally substituted with one or more substituents each independently selected from the group consisting of F, Cl, Br, NO<sub>2</sub>, amino,


N-alkylamino, N,N-dialkylamino, CN, optionally substituted alkyl, -O-(optionally substituted alkyl) and phenyl;

$Z^{110}$  and  $Z^{111}$  for each occurrence is independently ( $C_0-C_3$ ) optionally substituted with optionally substituted phenyl; and

5 A is -N(R)-C(O)-N(R)-, -N(R)-S(O)<sub>2</sub>-, -N(R)-C(O)-, -N(R)- or -N(R)-C(O)-O-.

80. A compound according to Claim 79, wherein  $R_4$  is methyl and there is only one  $R_a$  and it is F.

10 81. A compound according to Claim 9 or 66, wherein



$Z^{100}$  is an optionally substituted group selected from the group consisting of phenyl, isoxazolyl, tetrahydronaphthyl, furanyl, benzofuranyl, pyridyl and indolyl;


15 where  $Z^{100}$  is optionally substituted with one or more substituents each independently selected from the group consisting of F, CN, NO<sub>2</sub>, -C(O)H, -CONH<sub>2</sub>, -NHSO<sub>2</sub>CF<sub>3</sub>, optionally substituted alkyl, optionally substituted heteroaryl and -O-(optionally substituted alkyl);

$Z^{110}$  and  $Z^{111}$  are each independently optionally substituted ( $C_0-C_3$ ); and

A is O, -N(R)-C(O)-(CH<sub>2</sub>)<sub>n</sub>-N(R)-, -C(O)-N(R)-, -N(R)-C(O)-O-, -N(R)-C(O)- or -N(R)-.

20 82. A compound according to Claim 81, wherein  $R_4$  is methyl;  $R_a$  is H or methoxy; and  $Z^{110}$  and  $Z^{111}$  are each unsubstituted.


83. A compound according to Claim 9, wherein  $R_1$  is



where R is H or lower alkyl and n is for each occurrence is independently 1 to 6.


84. A compound according to Claim 83, wherein  $R_1$  is

5



85. A compound according to Claim 84, wherein  $Z^{100}$  is substituted or unsubstituted phenyl.

10 86. A compound according to Claim 8, 9 or 10, wherein



5    87. A compound according to Claim 11 wherein n is 2; R<sub>6</sub> is H; m is 1; r is 1; and R<sub>4</sub> and R<sub>5</sub> are each hydrogen.

88. A compound according to claim 64 or 87 wherein R<sub>1</sub> is 4-phenoxyphenyl.